

# UC Irvine

## UC Irvine Previously Published Works

### Title

Response to letter: contribution of serum ethanol concentration to the osmol gap: a prospective volunteer study.

### Permalink

<https://escholarship.org/uc/item/3jw9s0pv>

### Journal

Clinical toxicology (Philadelphia, Pa.), 51(8)

### ISSN

1556-3650

### Authors

Carstairs, SD  
Suchard, JR  
Smith, T  
et al.

### Publication Date

2013-09-01

### DOI

10.3109/15563650.2013.824584

Peer reviewed

LETTER TO THE EDITOR

**Response to Letter: Contribution of Serum Ethanol Concentration to the Osmol Gap: a Prospective Volunteer Study**

*To the Editor:*

We appreciate the comments by Sud et al. regarding the variability of the coefficient among our subjects and the utility of the osmol gap in determining whether a clinically significant toxic alcohol ingestion is present; indeed, we share some of these same concerns. However, our findings are in line with previous investigations on the subject. Various studies have shown that the osmol gap can vary significantly from person to person<sup>1</sup> and even within a single individual over time.<sup>2</sup>

The goal of our study was not to alter the coefficient used by clinicians in practice but rather to determine more accurately ethanol's contribution to the osmol gap in a controlled environment without the confounders found in previous studies. As Sud et al. pointed out, our coefficient of 4.25 is quite similar to the coefficient of 4.22 derived by Hoffman et al. and may represent a more accurate determination of the actual contribution of ethanol to serum osmolality. However, as we pointed out in the limitations section of our paper, the use of osmol gaps in the diagnosis of toxic alcohol poisoning has inherent limitations and should not be used as a substitute for good clinical decision-making. Fomepizole is now sold in generic form in the U.S. and is significantly less expensive than it has been in the past. If toxic alcohol ingestion is suspected, fomepizole should be administered empirically while awaiting confirmatory testing, and the decision to treat should not be based upon the presence or absence of an elevated osmol gap.

*S. D. Carstairs*

*Department of Emergency Medicine,  
Naval Medical Center, San Diego,  
CA, USA*

*J. R. Suchard*

*Department of Emergency Medicine,  
University of California, Irvine Medical Center,  
Orange, CA, USA*

*T. Smith, L. V. Simon & C. J. Kalynych*

*Department of Emergency Medicine,  
University of Florida College of Medicine,  
Jacksonville, FL, USA*

*M. Shimada*

*University of Tokyo, Tokyo, Japan*

*D. A. Tanen*

*Department of Emergency Medicine,  
Naval Medical Center,  
San Diego, CA USA*

**References**

1. Hoffman RS, Smilkstein MJ, Howland MA, Goldfrank LR. Osmol gaps revisited: normal values and limitations. *J Toxicol Clin Toxicol* 1993; 31:81–93.
2. Suchard JR. Intra-subject variability of the baseline serum osmolal gap [abstract]. *J Toxicol Clin Toxicol* 2005; 43:689–690.

Received 01 July 2013; accepted 8 July 2013.

Address correspondence to Dr. Shaun D. Carstairs, MD, Department of Emergency Medicine, Naval Medical Center San Diego, 34800 Bob Wilson Drive, San Diego, CA, 92134-5000. Email: shaunc@yahoo.com